題號: 185 國立臺灣大學 105 學年度碩士班招生考試試題

科目: 臨床牙周病學

題號: 185

共 2 頁之第 / 頁

節次: 4

## 第一大題 1-10,單選,每一小題2分,請將答案填寫於答案卷第一頁【選擇題作答區】內

- 1. In a healthy mandibular first molar, which type of cementum will be found mostly over furcation area?
- (A). Acellular afibrillar
- (B). Acellular extrinsic fiber
- (C). Cellular mixed stratified
- (D). Cellular intrinsic fiber
- 2. Which of the following structure is the specified source of junctional epithelium?
- (A). Reduced enamel epithelium
- (B). Oral gingival epithelium
- (C). Epithelial cell rests of Malassez
- (D). Hertwig's epithelial root sheath
- 3. The salivary MMP-8 (matrix metalloproteinase-8) level is correlated with chronic periodontitis, which of the following antibiotics may inhibit MMP-8?
- (A). Amoxicillin
- (B). Tetracycline
- (C). Metronidazole
- (D). Ciprofloxacin
- 4. Which periodontal bacteria will secret various virulent factors such as proteases, a collagenase, and a hemolysin?
- (A). A. actinomycetemcomitans
- (B). T. forsythia
- (C). P. gingivalis
- (D). T. denticola
- 5. Among oral flora, which of the following will be regarded as a true periodontal pathogen?
- (A). T. forsythia
- (B). C. rectus
- (C). F. nucleatum
- (D). P. intermedia
- 6. Which kind of the inflammatory cells is dorminated in the soft tissue of advanced periodontitis lesion?
- (A). neutrophils.
- (B). macrophages
- (C). plasma cells..
- (D). Lymphocytes
- 7. If you examine a periodontitis tooth with probe, where will the tip of probe usually stop at?

題號: 185 國立臺灣大學 105 學年度碩士班招生考試試題

科目:臨床牙周病學 題號: 185

節吹: 4 共 2 頁之第 2 頁

(A). Alveolar crest.

- (B). Connective tissue.
- (C). Apical portion of the junctional epithelium.
- (D). Coronal portion of the junctional epithelium.
- 8. Which of the following factor may most likely influence the probing depth in periodontally involved sites?
- (A). age of the patient.
- (B). thickness of the gingiva.
- (C). location of the probing site.
- (D). disease severity
- 9. R. C. Bower (1979) studied the furcation entrance diameter of the permanent molars and found what percent of all furcation sites were less than 0.75mm (the width of commonly used periodontal curettes)?
- (A). 18%
- (B). 38%
- (C). 58%
- (D). 78%
- 10. Which of the following sign or symptom is NOT commonly seen in patients with chronic periodontitis?
- (A). Papillary enlargement
- (B). Bone destruction.
- (C). Gingival bleeding.
- (D). Severe pain

## 第二大題 問答題:

- 11. Describe the role of buccal keratinized gingiva to the maintenance care of periodontal condition. (20%)
- 12. Why there will radiographic crestal bone loss be noticed around the implant, and how to prevent it? (20%)
- 13. How to choose an interdental cleaning aid, and describe the correct technique to use it. (20%)
- 14. Please describe the key factors which need be considered in determining the prognosis of a periodontal diseased tooth, and give your definitions about fair, questionable and hopeless prognosis. (20 %)

## 試題隨卷繳回