題號: 56

國立臺灣大學 106 學年度碩士班招生考試試題

科目:線性代數(A)

科日・緑性代数(A) 節次: 4 題號: 56 英 | 頁之第 | 頁

Notice: You must show all your work in order to receive full credit.

- (1) (20 points) Show that if $\mathbb{R}^n = W_1 \cup W_2 \cup \cdots \cup W_k \cup \cdots$, where each W_k is a subspace, then $\mathbb{R}^n = W_i$ holds for some i.
- (2) (20 points) Let a, b, c, d, e, f be real numbers such that the quadratic form $Q(x, y, z) := ax^2 + by^2 + cz^2 + 2dxy + 2eyz + 2fxz$ is positive definite. Then the region bounded by the surface Q(x, y, z) = 1 has volume equals

$$\frac{4\pi}{3\sqrt{abc+2def-ae^2-bf^2-cd^2}}.$$

- (3) (15 points) The are infinitely many t in \mathbb{R} such that the vectors $(t, 2t^2, 3t^3, 4t^4)$, $(t^2, 2t^3, 3t^4, 4)$, $(t^3, 2t^4, 3t, 4t^2)$, $(t^4, 2t, 3t^2, 4t^3)$ form a basis of \mathbb{R}^4 .
- (4) (15 points) Determine all values of $a,b,c,d,e,f\in\mathbb{R}$ such that the

$$\text{matrix } A := \begin{pmatrix} 1 & a & b & c \\ 0 & 1 & d & e \\ 0 & 0 & 2 & f \\ 0 & 0 & 0 & 2 \end{pmatrix} \text{ is } not \text{ diagonalizable.}$$

- (5) (10 points) If a 5×5 matrix $A \in M_5(\mathbb{R})$ satisfies $A^7 = I_5$ (the identity matrix), then 1 is an eigenvalue of A.
- (6) Prove or disprove the following statements (10 points for each).
 - (a) If $\psi: \mathbb{R}^n \longrightarrow \mathbb{R}^n$ is a linear transformation with the null space (kernel) of dimension n-1, then there exists some $v \in \mathbb{R}^n$ and a non-zero $\lambda \in \mathbb{R}$ such that $\psi(v) = \lambda \cdot v$.
 - (b) If W_1 and W_2 are 8-dimensional subspaces of \mathbb{R}^{10} , then there exist $a_1, a_2, \dots, a_{10}, b_1, b_2, \dots, b_{10}, c_1, c_2, \dots, c_{10}, d_1, d_2, \dots, d_{10}$ in \mathbb{R} such that the intersection $W_1 \cap W_2$ is the set of all vectors (x_1, \dots, x_{10}) with x_1, \dots, x_{10} a solution to the system of equations

$$\begin{cases} a_1x_1 + a_2x_2 + \dots + a_ix_i + \dots + a_{10}x_{10} &= 0 \\ b_1x_1 + b_2x_2 + \dots + b_ix_i + \dots + b_{10}x_{10} &= 0 \\ c_1x_1 + c_2x_2 + \dots + c_ix_i + \dots + c_{10}x_{10} &= 0 \\ d_1x_1 + d_2x_2 + \dots + d_ix_i + \dots + d_{10}x_{10} &= 0. \end{cases}$$

試題隨卷繳回