科目:遺傳學(B)

節次: 6

題號:450 共 7 頁之第 1 頁

一、選擇題 (每題2分,共50分)

1. When DNA is heated, the hydrogen bonds break and allow the two strands of DNA to
separate from one another. Which sequence of DNA would be most resistant to heating, and
therefore, stay double stranded longer?

1. When DNA is heated, the hydrogen bonds break and allow the two strands of DNA to
separate from one another. Which sequence of DNA would be most resistant to heating, an
therefore, stay double stranded longer?
a. AATTGGCC
b. ATGCATTC
c. AGGAGCTC
d. TGGCTTAG
e. TACCAATT
2. The individual credited with first observing the cell nucleus is
a. Anton van Leeuwenhoek
b. Robert Brown
c. Matthias Schleiden
d. Theodor Schwann
e. Rudolf Virchow
3. The tight packing of chromatin into the chromosomes visible during nuclear division is
possible due to links between proteins.
a. Cdk
b. H1 histone
c. H3 histone
d. H4 histone
e. nonhistone
4. If a cell contains 84 sister chromatids at metaphase of mitosis, how many chromosomes
will be present in each nucleus in late telophase?
a. 168
b. 84
c. 42

- d. 21
- e. cannot be determined from the information given

科目:遺傳學(B)

題號:450

節次: 6

共 7 頁之第 2 頁

5. Colchicine, a chemical extracted from the autumn crocus (Colchicum autumnale), inhibits
the formation of microtubules. Mitotic cells exposed to colchicine will therefore be arrested at
·
a. anaphase
b. interphase
c. prophase
d. telophase
e. metaphase
6. The microtubules making up the mitotic spindle attach to specialized structures called
that are found in the centromere region of the chromosome.
a. nucleosomes
b. centrosomes
c. kinetochores
d. chromatids
e. centrioles
7. During prokaryotic cell division, two chromosomes separate and are distributed to the two
ends of the cell by
a. the action of the mitotic spindle
b. an unknown mechanism
c. attachment to actin
d. attachment to separating membrane regions
e. formation of a newly made cell wall
8. Which statement describes a difference between meiosis I and meiosis II?
a. DNA is replicated during meiosis II but not during meiosis I.
<ul> <li>Homologous chromosomes separate during meiosis I; sister chromatids separate during meiosis II.</li> </ul>
c. Meiosis I produces diploid cells; meiosis II produces haploid cells.
d. Chromosomes align along the equatorial plane of the cell in meiosis I but not in meiosis II.
e. Crossing-over occurs during meiosis II but not during meiosis I.

題號: 450

## 國立臺灣大學110學年度碩士班招生考試試題

9. Homologous chromosomes undergo recombination during \_\_\_\_\_.

科目:遺傳學(B)

科日・通得学(B 節次: 6 題號:450

共 7 頁之第 3 頁

	a. prophase II		
	b. metaphase I		
	c. metaphase II		
d. both prophase I and II			
	e. prophase I		
	10. Nondisjunction during meiosis I results in		
	a. the generation of an extra daughter cell		
b. a change in the status of a daughter cell from diploid to haploid			
	c. a change in the status of a daughter cell from haploid to diploid		
	d. one pole of the cell receiving neither member of a homologous pair of		
	chromosomes		

- 11. If a species has 42 pairs of chromosomes, how many maternal and paternal chromosome combinations are possible during independent assortment?
  - a.  $42^2$
  - b. 2<sup>42</sup>
  - c. 2<sup>21</sup>
  - $d.21^{2}$
  - e. 84<sup>2</sup>
- 12. Assuming a single cross-over event, the completion of meiosis produces \_\_\_\_\_.
  - a. three daughter cells with parental chromosomes and one daughter cell with a recombinant chromosome
  - b. four daughter cells with recombinant chromosomes

e. a gamete that cannot fuse with another gamete

- c. two daughter cells with parental chromosomes and two daughter cells with recombinant chromosomes
- d. one daughter cell with parental chromosomes and one cell with recombinant chromosomes
- e. two daughter cells, each with recombinant chromosomes

科目:遺傳學(B)

題號: 450

節次: 6

共 7 頁之第 4 頁

13. Pai	rs of alleles are found at a particular on a pair of
a	gene; gametes
ь	base; homozygous genes
c	nucleotide; genes
đ	locus; homologous chromosomes
е	sister chromatid; homologous chromosomes
14. A	individual is always part of a testcross.
a	heterozygous
ь	homozygous recessive
С	homozygous dominant
đ	self-fertilized
е	dihybrid
	parent has a genotype of <i>RrYy</i> . What is the probability of this individual producing a with the <i>RY</i> genotype?
a	. 1/2
b	. 1/4
С	3/4
d	. 1/8
е	. 0
16. Y	our father is heterozygous for a recessive disorder. You know your mother has two
	" alleles. What is the probability that you will have the disorder?
a	. 0%
b	. 25%
c	. 50%
d	. 75%
e	. 100%

題號: 450

## 國立臺灣大學110學年度碩士班招生考試試題

科目:遺傳學(B)

行日 · 通符 節次 : 6 題號:450

共 7 頁之第 5 頁

17. Mouse pigmentation is subject to epistasis of the $B$ alleles by the $d$ alleles. $B$ (black) is
dominant over $b$ (brown), and $D$ is dominant over $d$ . Homozygous $d$ is epistatic to the black
and brown genes. Given this information, what genotype(s) result in a white mouse (no
pigment)?

- a. BBdd only
- b. Bbdd only
- c. bbDD only
- d. bbDd only
- e. BBdd and Bbdd

18. 2	Adjacent nucleotide	s on a strand of DNA	are connected to	each other by a	a(n)
-------	---------------------	----------------------	------------------	-----------------	------

- a. hydrophobic interaction
- b. phosphodiester bond
- c. hydrogen bond
- d. peptide bond
- e, ionic bond

10	Which nucleotide sequence	is complementary	to the DNA se	avence 5'-GACGTT-3'?
10	Which nucleotide sequence	e is complementary	to the DNA se	equence 3 -GACG11-3 (

- a. 5'-TCATGG-3'
- b. 3'-TCATGG-5'
- c. 3'-CTGCAA-5'
- d. 3'-AGTACC-5'
- e. 5'-TTGCAG-3'

#### 20. DNA polymerase \_\_\_\_\_.

- a. synthesizes a short RNA segment to begin DNA replication
- b. adds nucleotides to the 5' end of an existing strand to synthesize a new DNA strand
- c. adds nucleotides to the 3' end of an existing strand to synthesize a new DNA strand
- d. seals nicks between adjacent segments of DNA
- e. unwinds the DNA double helix at the origin of replication

科目:遺傳學(B)

題號:450

節次: 6

共 7 頁之第 6 頁

21. DNA replication is said to be semiconservative because						
a. the number of nucleotides within genes remains constant						
b. half of the DNA in a cell comes from one parent and the other half from the other						
parent.						
c. the same process of DNA replication is used by all organisms						
d. the total amount of DNA within an individual remains the same						
e. each new DNA molecule is composed of one old strand and one new strand						
e. each new DIVA molecule is composed of one old sidual and one new strand						
22. Reiji Okazaki discovered what are now called "Okazaki fragments" produced during						
DNA replication. These fragments are						
a. short lengths of new DNA on the leading strand						
b. RNA primers on the lagging strand						
c. RNA primers on both the lagging and leading strand						
d. short lengths of new DNA on the lagging strand						
e. RNA primers on the leading strand						
23. Individuals with xeroderma pigmentosum inherit a defective DNA repair mechanism. As						
a consequence,						
a. they are sterile						
b. their cells have no proofreading abilities during DNA replication						
c. Okazaki fragments produced during DNA replication cannot be joined						
d. they easily develop skin cancer when exposed to sunlight						
e. their telomeres are shorter than average						
24. The correction and a in gold to be degenerate because						
24. The genetic code is said to be degenerate because  a. some codons that do not specify an amino acid						
b. most amino acids are represented by more than one codon						
c. the code varies considerably between different organisms						
d. the code is commaless, with no indicators of spaces between codons						
e. the code varies considerably between different cell types within a multicellular						
organism						
· ·						
25. The TATA box is a key element of the of most eukaryotic protein-coding genes.						
a. terminator						
b. coding region						
c. promoter						
d. transcription start point						
e. introns						

科目:遺傳學(B)

節次: 6

#### 二、 問答題 (每題10分,共50分)

- Describe the roles of the following enzymes during DNA replication: ligase, singlestranded binding proteins (SSBs), DNA polymerase III, primase, DNA helicase.
- Although Watson and Crick are credited with the discovery of the structure of DNA, they
  based this discovery on the bench work of many other scientists of the time. Name at least
  two different pieces of knowledge that Watson and Crick used in their discovery that was
  based on work from other scientists.
- 3. Compare and contrast the cellular locations, templates, and products of transcription and translation in prokaryotes and eukaryotes.
- 4. A couple wants to start a family, but they are concerned that their child might have cystic fibrosis, which is a recessive trait.. After taking a family history, you determine that while neither of them have the disease, the woman had a sister with cystic fibrosis (with unaffected parents), and the man's father also had cystic fibrosis. What do you tell them?
- 5. Describe the meiotic events that generate genetic diversity in gametes.